

① Tree contributions are almost equal in the three different approaches.

⇒ almost same Branching ratio.

However the penguin contributions are larger in PQCD due to the dynamical enhancement!

$$\mu_{\text{Fact.}} : m_b \text{ or } m_b/2 \rightarrow \sqrt{\bar{\Lambda} m_b} \sim 1.56 \text{ GeV}$$

② Factorizable tree and Penguin contributions

- PQCD : real
- FA. BBNS : complex (BSS and FSI)

③ Important annihilation contribution!

- same order as the Factorized Penguin
- large imaginary part.
- the main source of strong phase!




Figure 9

④ Nonfactorizable contributions : M^P, M^T

- PQCD : complex
- BBNS : real
- FA : included in the Wilson Coeff.

⑤ T and P ? (unit: 10^{-3})

$T_{PQCD} = 83.40 + i 1.60 ; P_{PQCD} = -6.34 + i 3.89$

$T_{BBNS} = 74.93 + i 1.13 ; P_{BBNS} = 1.13 - i 1.27$

$T_{FA} = 75.70 + i 1.13 ; P_{FA} = -3.04 - i 1.27$

$\rightarrow |P_{PQCD}| \sim 4.4 |P_{BBNS}| \sim 2.3 |P_{FA}|$

⑥ Large strong phase of $B^0 \rightarrow \pi^+ \pi^-$;

$A = V_t^* |P| e^{i\delta} - V_u^* |T|$

• $\delta_{PQCD} = 147^\circ ; \left| \frac{P}{T} \right| = 27\%$

• $\delta_{BBNS} = -50^\circ$

• $\delta_{FA} = -113^\circ$


⑦ Large CP :

$|A_{CP}| < 30\% : PQCD$

$|A_{CP}| < 4\% : BBNS$

$|A_{CP}| < 10\% : FA$

CP asymmetry in $B^0 \rightarrow \pi^+\pi^-$.



Direct
 By measuring \surd CP asymmetry of $\pi^+\pi^-$,
 We can distinguish PQCD method from others!

B) B → Kπ decays:

21

TABLES

Amplitude	Real part	Imaginary part
F_e	103.69	0
F_c^P	-8.17	0
F_a^P	-0.23	3.75

TABLE I. Amplitudes for the $\bar{B}^0 \rightarrow K^- \pi^+$ decay in units of 10^{-2} with $F_e^P = F_{c4}^P + F_{c6}^P$ and $F_a^P = F_{a4}^P + F_{a6}^P$.

$\phi_3 = 80^\circ$

Branching ratio	PQCD prediction (10^{-6})		CLEO data (average) (10^{-6})
$\text{Br}(B^+ \rightarrow K^0 \pi^+)$	$17.51^{+3.3}_{-2.8}$	$16.99^{+3.3}_{-2.7}$	$18.2^{+4.6}_{-4.0} \pm 1.6$
$\text{Br}(B^- \rightarrow \bar{K}^0 \pi^-)$	$16.48^{+3.2}_{-2.7}$	$A_{cp} = (3.0^{+0.2}_{-0.1}) \%$	$18.2^{+4.6}_{-4.0} \pm 1.6$
$\text{Br}(B^0 \rightarrow K^+ \pi^-)$	$18.85^{+3.4}_{-2.7}$	$16.49^{+3.2}_{-2.5}$	$17.2^{+2.5}_{-2.4} \pm 1.2$
$\text{Br}(\bar{B}^0 \rightarrow K^- \pi^+)$	$14.13^{+3.0}_{-2.3}$	$A_{cp} = (-14.3^{+1.1}_{-1.0}) \%$	$17.2^{+2.5}_{-2.4} \pm 1.2$
$\text{Br}(B^\pm \rightarrow K^\pm \pi^0)$	$10.08^{+2.0}_{-1.6}$	$9.04^{+1.9}_{-1.5}$	$11.6^{+3.0+1.4}_{-2.7-1.3}$
$\text{Br}(B^\pm \rightarrow K^\mp \pi^0)$	$8.07^{+1.7}_{-1.4}$	$A_{cp} = (-11.5^{+0.9}_{-0.9}) \%$	$11.6^{+3.0+1.4}_{-2.7-1.3}$
$\text{Br}(B^0 \rightarrow K^0 \pi^0)$	$10.2^{+0.3}_{-0.3}$	$10.6^{+0.3}_{-0.3}$	$14.6^{+5.9+2.4}_{-5.1-3.3}$
$\text{Br}(\bar{B}^0 \rightarrow \bar{K}^0 \pi^0)$	$10.9^{+0.3}_{-0.2}$	$A_{cp} = (3.4^{+0.4}_{-0.3}) \%$	$14.6^{+5.9+2.4}_{-5.1-3.3}$

TABLE II. PQCD predictions of branching ratios for a representative parameters set.

$$\bullet \left| \frac{P}{T} \right| = \left| \frac{V_{ts}^* V_{tb}}{V_{us}^* V_{ub}} \right| \frac{(F_e^P + M_e^P + F_a^P + M_a^P)}{(F_e + M_e^T)}$$

~ 6.0

• Ratios of CP-averaged rates:

	PQCD	BBNS	CLEO
• $\frac{\text{Br}(\pi^+\pi^-)}{\text{Br}(\pi^\pm K^\mp)}$	= 0.45 ~ 0.80	0.5 ~ 1.9	[0.25 ± 0.10]

• $\frac{\text{Br}(\pi^\mp K^\pm)}{2 \text{Br}(\pi^0 K^0)}$	= 0.68 ~ 0.9	0.9 ~ 1.4	[0.59 ± 0.27]
---	--------------	-----------	---------------

• $\frac{2 \text{Br}(\pi^0 K^\pm)}{\text{Br}(\pi^\pm K^0)}$	= 1.06 ~ 1.08	0.9 ~ 1.3	[1.27 ± 0.47]
---	---------------	-----------	---------------

• $\frac{\tau_{B^+}}{\tau_{B^0}} \cdot \frac{\text{Br}(\pi^\mp K^\pm)}{\text{Br}(\pi^\pm K^0)}$	= 1.03 ~ 1.10	0.6 ~ 1.0	[1.00 ± 0.30]
---	---------------	-----------	---------------

• $\text{Br}(B \rightarrow \pi^0 K^0) = (10.6 \pm 0.3) \cdot 10^{-6}$: PQCD
 $= (4.5 \pm 2.5) \cdot 10^{-6}$: BBNS

Two-times larger !!


CP Asymmetry of $B^0 \rightarrow K^\pm \pi^\mp$ 

Figure 7




Figure 5(a)




Figure 5(b)




Figure 7(a)




Figure 7(c)

c) $B \rightarrow \phi K$ decay

detail will be in
Keum's poster.

- $B \rightarrow VP$ mode.
- Pure Penguin process.

• $B^{\pm} \rightarrow \phi K^{\pm}$:

$$A_{\nu} B_{r}(B^{\pm} \rightarrow \phi K^{\pm}) = (12.1^{+2.6}_{-1.9}) \cdot 10^{-6}$$

$$A_{cp} = 0.0\%$$

• $B^0 \rightarrow \phi \bar{K}^0$:

$$A_{\nu} B_{r}(B^0 \rightarrow \phi \bar{K}^0) = (9.0^{+2.5}_{-1.8}) \times 10^{-6}$$

$$A_{cp} = -(1.7^{+0.2}_{-0.1})\%$$

Summary:

With PQCD approach including t_{\perp} ,

1 We can calculate Form Factors (Space-like, timelike)



• Sizable t_{\perp} contributions

• Threshold Resummation

2 We have

Large Branching ratios
in charmless B decays
except $B \rightarrow \eta' K$



Dynamical
Penguin
Enhancement
at $\mu \sim 1.5 \text{ GeV}$

Large direct CP asym.
for $B \rightarrow \pi^+ \pi^-, K^+ \pi^-, K^+ \pi^0$



Large strong phase
from the factorized
Annihilation
diagram

BRs in units of 10^{-6}

Decay	CLEO	BELLE	BABAR
$K^{\pm}\pi^{\mp}$	$17.2^{+2.5}_{-2.4} \pm 1.2$	$17.4^{+5.1}_{-4.6} \pm 3.4$	$12.5^{+3.0+1.3}_{-2.6-1.7}$
$K^0\pi^{\pm}$	$18.2^{+4.6}_{-4.0} \pm 1.6$	$16.6^{+9.8+2.2}_{-7.8-2.4}$	
$K^{\pm}\pi^0$	$11.6^{+3.0+1.4}_{-2.7-1.3}$	$18.8^{+5.5}_{-4.9} \pm 2.3$	
$K^0\pi^0$	$14.6^{+5.9+2.4}_{-5.1-3.3}$	$21.0^{+9.3+2.5}_{-7.8-2.3}$	
$\pi^{\pm}\pi^{\mp}$	$4.3^{+1.6}_{-1.4} \pm 0.5$	$6.3^{+3.9}_{-3.5} \pm 1.6$	$9.3^{+2.6+1.2}_{-2.3-1.4}$
$\pi^{\pm}\pi^0$	< 12.7	< 10.1	
$\pi^0\pi^0$	< 5.7		
$K^{\pm}\eta'$	$80^{+10}_{-9} \pm 7$		$62 \pm 18 \pm 8$
$K^0\eta'$	$89^{+18}_{-16} \pm 9$		
$*\rho^{\pm}\pi^{\mp}$	$27.6^{+8.4}_{-7.4} \pm 4.2$		$49 \pm 13^{+6}_{-5}$
$\rho^0\pi^{\pm}$	$10.4^{+3.3}_{-3.4} \pm 2.1$		
$\omega\pi^{\pm}$	$11.3^{+3.3}_{-2.9} \pm 1.4$		
$K^{*\pm}\pi^{\mp}$	22^{+8+4}_{-6-5}		
$K^{*0}\eta$	$13.8^{+5.5}_{-4.6} \pm 1.6$		
$K^{*\pm}\eta$	$26.4^{+9.6}_{-8.2} \pm 3.3$		
ϕK^{\pm}	$6.4^{+2.5+0.5}_{-2.1-2.0}$	$17.2^{+6.7+1.8}_{-5.4-1.8}$	

PQCD

$$16.5^{+3.2}_{-2.5}$$

$$17.0^{+3.3}_{-2.7}$$

$$9.1^{+1.9}_{-1.5}$$

$$10.6 \pm 0.3$$

$$8.2^{+2.0}_{-1.5}$$

$$\sim 3.7^*$$

$$\sim 0.2^*$$

$$\sim 18.4^*$$

$$\sim 34.0^*$$

$$\sim 8.0^*$$

$$12.1^{+2.6}_{-1.9}$$

$$9.0^{+2.5}_{-1.8}$$

ϕK^0

* K. Ukai

* E. Kou

* C. D. Lü, M. Yang

□ Our results is well agreed with
experimental data for
Nonleptonic charmless B-meson decays.

→ See Table !

To PQCD Group members,
Thanks for your collaboration
and contributions !

E. Kou

T. Kurimoto

Hsiang-nan Li, Chun-Hung Chen

C. D. Lü, M. Nagashima

T. Morozumi, S. Mishima

N. Sinha, R. Sinha

K. Ukai

T. Yoshikawa

M. Z. Yang

A. I. Sanda